Author index

Anderton, M., see Canet-Aviles, R.-M. (102) 62

Atkins, H.D., see Thompson, C.L. (102) 55

Brown, E.M., see Chattopadhyay, N. (102) 73

Cai, D., see Yu, E.Z. (102) 9

Canet-Aviles, R.-M., Anderton, M., Hooper, N.M., Turner, A.J. and Vaughan, P.F.T. Muscarine enhances soluble amyloid precursor protein secretion in human neuroblastoma SH-SY5Y by a pathway dependent on protein kinase C_{α} , src-tyrosine kinase and extracellular signal-regulated kinase but not phospholipase C (102) 62

Chattopadhyay, N., Tfelt-Hansen, J. and Brown, E.M. PKC, p42/44 MAPK and p38 MAPK regulate hepatocyte growth factor secretion from human astrocytoma

secretion from human astrocytoma cells (102) 73

Chazot, P.L., see Thompson, C.L. (102) 55 Chen, Q., see Ma, E. (102) 100

Chrétien, M., see Mbikay, M. (102) 35
Cristobal, R., Popper, P., Lopez, I., Micevych, P., De Vellis, J. and Honrubia, V.
In vivo and in vitro localization of brain-derived neurotrophic factor, fibroblast growth factor-2 and their receptors in the bullfrog vestibular end organs (102) 83

De Vellis, J., see Cristobal, R. (102) 83 Drewery, D.L., see Thompson, C.L. (102) 55

Givalois, L., Li, S. and Pelletier, G.

Central nitric oxide regulation of the hypothalamic-pituitary-adrenocortical axis in adult male rats (102) 1

Gulya, K., see Kovacs, B. (102) 28

Haddad, G.G., see Ma, E. (102) 100 Hallenbeck, J.M., see Yu, E.Z. (102) 9 Heike, T., see Ishibashi, H. (102) 110

Hihara, S., see Ishibashi, H. (102) 110 Hollingsworth, Z.R., see Spektor, B.S. (102)

Honma, M., see Iwata, T. (102) 105 Honrubia, V., see Cristobal, R. (102) 83 Hooper, N.M., see Canet-Aviles, R.-M. (102)

Iriki, A., see Ishibashi, H. (102) 110

Ishibashi, H., Hihara, S., Takahashi, M., Heike, T., Yokota, T. and Iriki, A. Tool-use learning induces BDNF expression in a selective portion of monkey anterior parietal cortex (102)

Iwata, T., Namikawa, K., Honma, M., Mori, N., Yachiku, S. and Kiyama, H. Increased expression of mRNAs for microtubule disassembly molecules during nerve regeneration (102) 105

Johnson, J.M., see Spektor, B.S. (102) 118

Kalenga, L., see Mbikay, M. (102) 35 Kaneko, Y.A., see Spektor, B.S. (102) 118 Kiyama, H., see Iwata, T. (102) 105 Kovacs, B. and Gulya, K. Differential expression of multiple

Differential expression of multiple calmodulin genes in cells of the white matter of the rat spinal cord (102) 28

Li, S., see Givalois, L. (102) 1 Liu, Y., see Wilson, M.E. (102) 48 Lopez, I., see Cristobal, R. (102) 83 Luthi-Carter, R., see Spektor, B.S. (102) 118

Ma, E., Tucker, M.C., Chen, Q. and Haddad, G.G.
 Developmental expression and enzymatic activity of pre-mRNA deaminase in *Drosophila* melanogaster (102) 100

Mbikay, M., Raffin-Sanson, M.-L., Sirois, F., Kalenga, L., Chrétien, M. and Seidah, N.G.

Characterization of a repressor element in the promoter region of proprotein convertase 2 (PC2) gene (102) 35

McCarron, R.M., see Yu, E.Z. (102) 9 Micevych, P., see Cristobal, R. (102) 83 Miller, D.W., see Spektor, B.S. (102) 118 Mishina, M., see Sakamoto, T. (102) 113 Mori, N., see Iwata, T. (102) 105

Namikawa, K., see Iwata, T. (102) 105 Niki, H., see Sakamoto, T. (102) 113

Ozaita, A. and Vega-Saenz de Miera, E. Cloning of two transcripts, HKT4.1a and HKT4.1b, from the human two-pore K⁺ channel gene KCNK4. Chromosomal localization, tissue

distribution and functional expression (102) 18

Pelletier, G., see Givalois, L. (102) 1 Penney, J.B., see Spektor, B.S. (102) 118 Popper, P., see Cristobal, R. (102) 83

Raffin-Sanson, M.-L., see Mbikay, M. (102)

Sakamoto, T., Mishina, M. and Niki, H. Mutation of NMDA receptor subunit epsilon 1: effects on audiogenic-like seizures induced by electrical stimulation of the inferior colliculus in mice (102) 113

Seidah, N.G., see Mbikay, M. (102) 35
Sirois, F., see Mbikay, M. (102) 35
Solano, S.M., see Spektor, B.S. (102) 118
Spektor, B.S., Miller, D.W., Hollingsworth,
Z.R., Kaneko, Y.A., Solano, S.M.,
Johnson, J.M., Penney, J.B., Young, A.B.
and Luthi-Carter, R.
Differential D₁ and D₂ receptormediated effects on immediate early gene induction in a transgenic mouse

Stephenson, F.A., see Thompson, C.L. (102)

model of Huntington's disease (102)

Takahashi, M., see Ishibashi, H. (102) 110 Tfelt-Hansen, J., see Chattopadhyay, N. (102)

Thompson, C.L., Drewery, D.L., Atkins, H.D., Stephenson, F.A. and Chazot, P.L. Immunohistochemical localization of *N*-methyl-D-aspartate receptor subunits in the adult murine hippocampal formation: evidence for a unique role of the NR2D subunit (102) 55

Tucker, M.C., see Ma, E. (102) 100 Turner, A.J., see Canet-Aviles, R.-M. (102) 62

Vaughan, P.F.T., see Canet-Aviles, R.-M. (102) 62

Vega-Saenz de Miera, E., see Ozaita, A. (102)

Wilson, M.E., Liu, Y. and Wise, P.M. Estradiol enhances Akt activation in cortical explant cultures following neuronal injury (102) 48 Wise, P.M., see Wilson, M.E. (102) 48

Yachiku, S., see Iwata, T. (102) 105 Yokota, T., see Ishibashi, H. (102) 110 Young, A.B., see Spektor, B.S. (102) 118 Yu, E.Z., Hallenbeck, J.M., Cai, D. and McCarron, R.M. Elevated arylalkylamine-N- acetyltransferase (AA-NAT) gene expression in medial habenular and suprachiasmatic nuclei of hibernating ground squirrels (102) 9